

Dirichlet String Theory and Singular Random Surfaces

Miao Li¹

Department of Physics
Brown University
Providence, RI 02912

We show that string theory with Dirichlet boundaries is equivalent to string theory containing surfaces with certain singular points. Surface curvature is singular at these points. A singular point is resolved in conformal coordinates to a circle with Dirichlet boundary conditions. We also show that moduli parameters of singular surfaces coincide with those of smooth surfaces with boundaries. Singular surfaces with saddle points indeed arise in the strong coupling expansion in lattice QCD. The kind of saddle point, which may be the origin of a singular point we need, is of infinite order.

¹ E-mail: li@het.brown.edu

Green proposed to study string theory propagating with surfaces bordered by Dirichlet boundaries, in order to reproduce the high energy behavior of scattering amplitudes in QCD [1]. Earlier references in the subject include those in [2]. For a recent approach to this string theory, see [3]. Is there any evidence based upon more theoretical grounds in support of the Dirichlet string theory, other than merely requiring account for “phenomenological” high energy results? The purpose of this note is to provide such evidence.

Starting with QCD, either from ’t Hooft’s $1/N$ expansion [4] or from Wilson’s strong coupling expansion on the lattice [5], one arrives at the conclusion that QCD may be described by a non-interacting string theory at the large N limit. Unfortunately this turned out not quite to be the case [6], at least some modifications are needed. Kazakov proposed a modified strong coupling expansion in [7], in which not only more degrees of freedom are required, but also some singular saddle points appear on random surfaces. This approach was further developed in [8], for a recent review, see [9]. Despite subtleties in going from the strong coupling regime to the weak coupling regime, those saddle points nevertheless survive. It is these singular points which interest us here. We shall argue that these points could serve as the origin of Dirichlet boundaries.

In the strong coupling expansion, a saddle point arises when $2n$ ($n > 1$) plaquettes are contracted cyclically on a link. The deficit angle at this point is $2\pi(n - 1)$. If this point survives the continuum limit, this deficit angle will induce a singular curvature, i.e., the curvature blows up approaching this point. Associated to a saddle point of a given order, there is a coefficient depending on the type of the saddle point. For the simplest saddle point with $n = 2$, there is an additional weight factor $1/N$. There are higher order corrections in terms of $1/N$. These can be explained as the coalescing of several saddle points of the same order. Now let us assume that this singularity can be described by a singular intrinsic curvature on the continuum surface. For a finite n , the curvature behaves as a δ -function centering at the saddle point. This kind of singularity will turn out to be too mild to induce a Dirichlet boundary. Nevertheless let us discuss this singularity first. Note that a saddle point of finite order is precisely a branch point of the same order found in the QCD_2 string theory [10]. To see this, consider a metric $ds^2 = |z|^{2(n-1)}|dz|^2$. The curvature of this metric behaves as $\sqrt{g}R \sim \delta^2(z)$. Use coordinates $w = z^n$ instead, the metric becomes smooth $ds^2 = |dw|^2$. The singular property is encoded in the fact that the z plane covers the w plane n times, and $z = 0$ is a branch point of order n . It is instructive to define the deficit angle by

$$\frac{\int_C ds}{\int_0^r ds} - 2\pi, \quad (1)$$

where C is the contour with a constant radius r . The first term is just the ratio of the circumference to the radius. In polar coordinates $z = re^{i\phi}$, the metric reads $ds^2 = r^{2(n-1)}(r^2 d\phi^2 + dr^2)$. It is easy to check that our definition of the deficit angle yields a value $2\pi(n-1)$ in this case. To study an appropriate conformal field theory, which is needed in a string theory, on a closed surface with such a branch point, one may either resolve this mild singularity by going to a higher genus (as done in the QCD₂ string theory), or one may simply insert an appropriate vertex operator [11] at this point and deal with a smooth surface (in coordinates w).

Now the kind of singular point inducing a Dirichlet boundary is different from a saddle point of any finite order. The reason is that the curvature blows up badly at the singular point. Without loss of generality, let us consider a metric

$$ds^2 = r^{2a} d\phi^2 + dr^2. \quad (2)$$

This metric is singular at $r = 0$ when $a < 1$. We also require $a > 0$, otherwise $r = 0$ is not really a point. A simple calculation yields $R = 2a(1-a)r^{-2}$. This curvature is so singular that the integral $\int \sqrt{g}R$ diverges at $r = 0$. Note that we have to put the restriction $a < 1$, otherwise $\int \sqrt{g}R$ converges. Thus, this singular point can not be resolved by going to higher genus. This is not surprising, since the deficit angle as defined in (1) is $2\pi(r^{a-1} - 1)$, which approaches infinity as r approaches zero. If we are to explain this singularity as originating from saddle points, it must consist of infinitely many saddle points on the same link with arbitrary large order. We conjecture that this is the case, although it is hard to have a derivation directly from the strong coupling expansion. After all the strong coupling expansion has not been proven to have a continuum limit yet.

We shall not assume any particular conformal field theory (CFT) in the QCD string theory here. To study a CFT with geometry (2), it is necessary to find a coordinate system in which the metric reads $\exp(\phi)|dz|^2$. Then we need to study this CFT in the geometry $|dz|^2$ by a Weyl rescaling. It is straightforward to see that the metric (2) can be written as $r^{2a}|dw|^2$, where $w = \frac{1}{1-a}r^{1-a} + i\phi$. Now w furnishes complex coordinates on the semi-infinite cylinder, and the boundary of this cylinder is at $r = 0$. Notice that if $a = 1$, one can not use these coordinates. The semi-infinite cylinder can be further mapped to the complex plane with a unit disk removed, by $z = \exp(w)$. The metric reads in these coordinates

$$ds^2 = \frac{1}{|z|^2} \left(\frac{1-a}{2} \ln|z|^2 \right)^{\frac{2a}{1-a}} |dz|^2.$$

The Weyl factor $\exp(\phi)$ vanishes on the boundary $|z| = 1$. The Liouville action, resulting from a Weyl rescaling, is infinite with this Weyl factor. We have seen that in a conformal coordinate system, the singular point *necessarily* blows up to a circle. If the CFT to be studied includes spacetime coordinates X^μ as a subset of conformal fields, the boundary conditions for these fields are Dirichlet, as this boundary represents just a previous singular point of curvature. We also note that the resulting conformal geometry is independent of parameter a . Formally, the Liouville action depends on a , however, we shall assume that this Liouville action be eventually canceled out by that from contribution of b - c ghosts. We postpone discussion of boundary conditions for these ghosts to a little later.

The above consideration of resolving a singular point can be generalized to the sphere with a singular point, and in general to a sphere with many such singular points. An interesting question then arises. If any such singular geometry is resolved resulting in a sphere with boundaries, are there enough configurations of singular geometries to cover the moduli space of the conformal sphere with boundaries? The answer to this question is positive. We consider the sphere with two singular points as an example. Assume that $r = 0$ and $r = \infty$ be two singular points on the sphere, in polar coordinates. The metric is approximately as in (2) around $r = 0$, with parameter a_1 . The metric is approximately $r'^{2a_2} d\phi^2 + dr'^2$ around $r = 1/r' = \infty$. a_i are all positive and less than one. We can write down such a metric readily

$$ds^2 = (1 + r^2)^{-2} ((r^{2a_1} + r^{4-2a_2})d\phi^2 + dr^2), \quad (3)$$

r^{2a_1} dominates r^{4-2a_2} when $r \rightarrow 0$; while r^{4-2a_2} dominates r^{2a_1} when $r \rightarrow \infty$. The above metric is written in conformal coordinates as

$$ds^2 = (1 + r^2)^{-2} (r^{2a_1} + r^{4-2a_2}) \frac{|dz|^2}{|z|^2}, \quad (4)$$

where

$$z = e^{R+i\phi}, \quad R(r) = \int_0^r \frac{dr}{(r^{2a_1} + r^{4-2a_2})^{1/2}}.$$

Apparently R is a well-defined function of r . The singular point $r = 0$ is mapped to $|z| = 1$, the unit circle. The singular point $r = \infty$ is mapped to the circle $|z| = \exp(R_0) > 1$. R_0 is finite, and given by

$$R_0 = \int_0^\infty \frac{dr}{(r^{2a_1} + r^{4-2a_2})^{1/2}} = \frac{1}{\sqrt{\pi}(4 - 2a_1 - 2a_2)} \Gamma\left(\frac{1 - a_1}{4 - 2a_1 - 2a_2}\right) \Gamma\left(\frac{1 - a_2}{4 - 2a_1 - 2a_2}\right). \quad (5)$$

We thus see that the sphere with two singular points is mapped to a conformal sphere with two disks removed, and it is just the annulus. The single real moduli parameter is the radius of the outer boundary $\exp(R_0)$, which is finite. The moduli parameter depends on both a_1 and a_2 . Adjusting a_i , R_0 can be sent to infinity. However, R_0 can never be zero, so geometries as described by (3) can never cover the whole moduli space. To cover the whole moduli space, one can replace the factor $r^{2a_1} + r^{4-2a_2}$ in (3) by $(r^{2a_1} + r^{4-2a_2})\rho^2(r)$. Here $\rho(r)$ is a function with $\rho(0) = \rho(\infty) = 1$. Now R_0 is not determined by (5) but by

$$R_0 = \int_0^\infty \frac{dr}{(r^{2a_1} + r^{4-2a_2})^{1/2} \rho(r)}.$$

It is possible to adjust $\rho(r)$ to let R_0 approach zero. This is done by increasing $\rho(r)$ for all r except $r = 0, \infty$. Geometrically, the result of doing this is that the circumference of a circle with constant r is increased. Notice that the distance between two points $r = 0$ and $r = \infty$ remains the same. This picture then explains why in conformal geometry, the outer boundary becomes closer and closer to the inner boundary. The whole moduli space is covered in this way. The moduli parameter then depends not only on parameters a_i , but also on details of geometry. However, the existence of boundaries is due to the existence of singular points.

These considerations generalize to a sphere with many singular points. It is hard and unnecessary to work out details of mapping singular geometries to conformal geometries with boundaries. We shall only argue that the whole moduli space of the latter is covered by varying singular geometries. Suppose there are $N \geq 3$ singular points, therefore there are N parameters a_i characterizing these singularities. Together with the positions of these singularities, there are $3N$ real parameters available. Since there is a global conformal invariance group $SL(2, C)$ on the sphere, only $2N - 6$ parameters out of $2N$ position parameters are relevant. So we are left with $3N - 6$ real parameters. This number is precisely the dimension of the moduli space of the sphere with N components of boundary. As in the $N = 2$ case, moduli parameters are complicated functions of the parameters of singular geometries, and it may be impossible to cover the whole moduli space if we consider only one class of geometries, such as that in (3). The moduli space can be covered by considering all possible singular geometries, again as in the $N = 2$ case. We believe that the conclusion presented in this section applies equally to Riemann surfaces of higher genus with singular points.

After having shown that Dirichlet string theory is equivalent to a closed string theory with singular surfaces, we come to the issue of b - c ghosts. If the boundary conditions for

scalar fields X^μ are Dirichlet, what are the boundary conditions for b - c ? Apparently, we can not impose Dirichlet boundary conditions on b - c consistently. These fields are fermionic, so Dirichlet boundary conditions can either be $b = 0$ or $c = 0$ on the boundary. If $b = 0$, then there is no nontrivial solution to the equation of motion $\bar{\partial}b = 0$, this would contradict the existence of moduli parameters. If $c = 0$, the trivial solution to $\bar{\partial}c = 0$ contradicts the existence of nontrivial conformal Killing vectors on, say, the disk. So we are left only with the possibility of Neumann type boundary conditions, namely $c\partial = \bar{c}\bar{\partial}$ and $b(dz)^2 = \bar{b}(d\bar{z})^2$ on the boundary. We now show how this is consistent with consideration of singular points. Consider the ghost fields in the geometry (2). For the c field, there are two components (c^r, c^ϕ) in polar coordinates. In conformal coordinates $(z, \bar{z})$, the two components become $c^z = zr^{-a}c^r + izc^\phi$ and $c^{\bar{z}} = \bar{z}r^{-a}c^r - i\bar{z}c^\phi$. So both c^z and $c^{\bar{z}}$ become singular on the boundary, if $c^r \neq 0$. If we demand that c^r approach zero faster than r^a (which approaches zero more slowly than r), we find the desirable Neumann boundary conditions. As for the b field, we find that two terms $b_{rr}(\partial r/\partial z)^2$ and $b_{r\phi}(\partial r/\partial z)(\partial\phi/\partial z)$ in b_{zz} are vanishing when $r \rightarrow 0$, for $\partial r/\partial z$ approaches zero. Neumann boundary conditions for this field then follow.

We showed in [3] that the Liouville action resulting from a Dirichlet boundary is essentially the same as the one from a Neumann boundary. In other words, Dirichlet boundary conditions do not change the central charge of the CFT. With a b - c ghost system obeying Neumann boundary conditions, it is possible to cancel the Liouville action from the matter CFT part. Together with our discussion above on the singular surfaces, we conclude that study of a string theory with such surfaces is made possible by studying the Dirichlet string theory. Because the latter makes perfect sense, it is then a reasonable hypothesis that singular surfaces contribute significantly in certain circumstances, such as in the QCD string theory.

As argued by Green, as well as in [3] for a general case, the Dirichlet boundaries provide us with a mechanism to generate the appropriate high energy behavior observed in strong interactions, which an ordinary string theory fails to do. Why is this so from the standpoint of the strong coupling expansion? The rest of this note is devoted to a plausible answer to this question.

First, let us remind ourselves how scattering amplitudes at high energies can be calculated in the pure glue theory. Color singlet states are supposed to be glueballs. A glueball state, as an eigen-state of four momentum, is a superposition of many states with different numbers of gluons. Determination of the wave function of a particular glueball state

is a nonperturbative problem. Fortunately, as a high energy process at a wide angle is concerned, there must be a factorization theorem, similar to the one for meson scattering [12]. This theorem states that the scattering amplitude factorizes into two factors. One of them has to do with the structure of the glueball, which may be called the distribution amplitude; another is the scattering amplitude between two individual gluons. The latter can be calculated perturbatively. Even the dependence of the distribution amplitude on the momentum transfer can be calculated perturbatively. The dependence of the whole amplitude on the large momentum transfer is largely determined by the two gluon scattering amplitude. This is translated into the short distance behavior of the two point function of the gauge potential $\langle A_\mu(x_1)A_\nu(x_2) \rangle$, which is calculated perturbatively.

Instead of considering high energy scattering amplitudes of glueballs, one can consider correlation functions of gauge invariant local operators constructed from field strength. High energy behavior is translated into certain short distance behavior of these correlation functions. As far as any short distance behavior is concerned, one can use the operator product expansion, again calculated perturbatively [13]

In the modified strong coupling expansion of Kazakov [7], a Lagrangian multiplier α , a hermitian matrix, is introduced to enforce unitarity of the link variable. Weight of each saddle point is the expectation value of a negative momentum of α , namely $W_n = \langle \frac{1}{N} \text{tr} \alpha^{-2n} \rangle$, where n is the order of the saddle point. It can be shown in the one plaquette model that solutions of the weight exist in both the strong coupling phase and the weak coupling phase [9]. We are certainly interested in the weak coupling phase. Here one finds that for large β , the inverse of the coupling constant, W_n goes to β^{-2n} . This factor cancels the factor β^{2n} from area of $2n$ plaquettes contracted on the link. This shows that all saddle points of higher order contribute in the weak coupling phase.

Similar arguments lead to demonstration of the existence of the weak coupling phase in QCD₄ [9], the physical model we are mostly interested in. Curiously, Kazakov showed that the weight W_n is associated to local condensation of the gluon field. Absorbing a factor β^{2n} into the weight, he finds

$$W_n = \beta \langle \frac{1}{N} \text{tr} F_{\mu\nu}^2 \rangle + \dots \quad (6)$$

The quantity on the r.h.s. of (6) can be calculated perturbatively in the weak coupling regime, using the standard Feynman rules. Now consider correlation functions of gauge invariant local operators which in turn are constructed from plaquette variables, in the

framework of strong coupling expansion. Saddle points appear in any such correlation function, and the weight in the weak coupling phase is given by eq.(6). When two operators in the correlation approach each other, we expect that saddle points near these operators play a more and more important role. It is then expected that the operator product expansion in the lattice context is dominated by saddle points, this is consistent with the fact that the weight in eq.(6) is determined perturbatively.

A lesson about the underlying CFT in the QCD string theory is readily drawn from the above discussion. In addition to the power-law behavior in the coefficients of the operator product expansion, there are logarithmic corrections, as functions of $\ln(p^2/\Lambda_{QCD}^2)$. The QCD scale Λ_{QCD} is the only free parameter in QCD. This parameter is replaced by the string tension $T \sim (\alpha')^{-1}$ in the string theory. So it is expected that similar factors involving $\ln(p^2/T)$ appear in the string calculations, even when only one singular point on the surface is present. With the choice of the simplest CFT described by free fields X^μ on the world sheet, it is shown in [1] and [3] that no such logarithmic factors are obtained. Thus, some modified CFT is required to yield logarithmic corrections. This is in contrast to the belief that logarithmic corrections arise from renormalizing infinities in the string theory. If this were true, one would be forced to introduce another independent scale. It is questionable that such a scale can be related to the string tension in any reasonable manner. From the strong coupling expansion, it is obvious that the free CFT is not the right theory, for example back-tracking configurations are not suppressed in this theory. It is likely that the correct CFT for the QCD string, if any, must involve interacting fields X^μ and additional fermionic fields to give logarithmic corrections as well as to suppress back-tracking.

In conclusion, no matter what the CFT may look like in the QCD string theory, singular surfaces must be taken into account in order to produce power-law behavior in high energy scattering amplitudes. We have shown that singular points can be resolved by introducing Dirichlet boundaries, so the theory can be studied systematically.

Acknowledgments

We would like to thank C.-I Tan for spiritual support, and for asking the question if it is possible to understand the heterotic pomeron in some string theory. We would also like to thank B. Urosevic, W. Zhao for comments, and A.T. Sornborger for reading the manuscript. This work was supported by DOE grant DE-FG02-91ER40688-Task A.

References

- [1] M.B. Green, Phys. Lett. B201 (1988) 42; B266 (1991) 325; B282 (1992) 380.
- [2] J.H. Schwarz, Nucl. Phys. B65 (1973) 131; E. Corrigan and D. Fairlie, Nucl. Phys. B91 (1975) 527; M. Green, Nucl. Phys. B103 (1976) 333.
- [3] M. Li, Dirichlet strings, BROWN-HET-915, hep-th/9307122.
- [4] G. 't Hooft, Nucl. Phys. B72 (1974) 461.
- [5] K. Wilson, Phys. Rev. D8 (1974) 2445.
- [6] D. Weingarten, Phys. Lett. B90 (1980) 285.
- [7] V. Kazakov, Phys. Lett. B128 (1983) 316; JETP 85 (1983) 1887.
- [8] I. Kostov, Phys. Lett. B138 (1984) 191; B147 (1984)445; K.H. O'Brien and J.-B. Zuber, Nucl. Phys. B253 (1985) 621; Phys. Lett. B144 (1984) 407.
- [9] V. Kazakov, Lectures given in the Trieste Spring School and Workshop-1993 on String Theory, hep-th/9308135.
- [10] D.J. Gross and W. Taylor, Nucl. Phys. B400 (1993) 181; B403 (1993) 395; J. Minahan, Phys. Rev. D47 (1993) 3430.
- [11] see for example, V.G. Knizhnik, Commun. Math. Phys. 112 (1987) 567.
- [12] see articles in Perturbative Quantum Chromodynamics, ed. A.H. Mueller, World Scientific, 1989.
- [13] H.D. Politzer, Phys. Rep. 14C (1974) 131.